

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## 4,4'-(1,8-Naphthalene-1,8-diyl)-dibenzonitrile

Carlos F. Lima,<sup>a</sup> Ligia R. Gomes,<sup>b</sup> Luís M. N. B. F. Santos<sup>a</sup> and John Nicolson Low<sup>c\*</sup>

<sup>a</sup>Centro de Investigação em Química, Departamento de Química e Bioquímica, Faculdade de Ciências, Universidade do Porto, Rua do Campo Alegre, 687, P-4169 007 Porto, Portugal, <sup>b</sup>REQUIMTE, Departamento de Química e Bioquímica, Faculdade de Ciências, Universidade do Porto, Rua do Campo Alegre, 687, P-4169 007 Porto, Portugal, and <sup>c</sup>Department of Chemistry, University of Aberdeen, Meston Walk, Old Aberdeen AB24 3UE, Scotland  
Correspondence e-mail: jnlow111@gmail.com


Received 29 November 2010; accepted 3 December 2010

Key indicators: single-crystal X-ray study;  $T = 150$  K; mean  $\sigma(\text{C}-\text{C}) = 0.002$  Å;  $R$  factor = 0.048;  $wR$  factor = 0.131; data-to-parameter ratio = 19.7.

In the title molecule,  $\text{C}_{24}\text{H}_{14}\text{N}_2$ , the exterior C—C—C angle of the naphthalene ring system involving the two phenyl-substituted C atoms is  $126.06$  (11)° and the dihedral angles between the mean plane of the naphthalene ring system and those of the benzene rings are  $66.63$  (5) and  $67.89$  (5)°. In the crystal, molecules are linked into a ladders by four weak C—H $\cdots\pi$  interactions.

## Related literature

For the structure of the related compound 4-(1-naphthyl)benzonitrile, see: Lima *et al.* (2010).



## Experimental

## Crystal data

$\text{C}_{24}\text{H}_{14}\text{N}_2$   
 $M_r = 330.37$   
 Monoclinic,  $C2/c$   
 $a = 17.0872$  (9) Å  
 $b = 8.2997$  (4) Å  
 $c = 24.3656$  (13) Å  
 $\beta = 93.795$  (2)°  
 $V = 3447.9$  (3) Å<sup>3</sup>  
 $Z = 8$   
 Mo  $K\alpha$  radiation  
 $\mu = 0.08$  mm<sup>-1</sup>  
 $T = 150$  K  
 $0.40 \times 0.30 \times 0.02$  mm

## Data collection

Bruker SMART APEX diffractometer  
 Absorption correction: multi-scan (SADABS; Bruker, 2004)  
 $T_{\min} = 0.971$ ,  $T_{\max} = 0.999$   
 11422 measured reflections  
 4634 independent reflections  
 3482 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.031$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.048$   
 $wR(F^2) = 0.131$   
 $S = 1.04$   
 4634 reflections  
 235 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.38$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.26$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

Cg1 and Cg2 are the centroids of the C1—C10 and C8—C10 rings, respectively.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C12}-\text{H12}\cdots\text{Cg2}^{\text{i}}$	0.95	2.75	3.6147 (13)	152
$\text{C16}-\text{H16}\cdots\text{Cg2}^{\text{ii}}$	0.95	2.92	3.6539 (15)	135
$\text{C82}-\text{H82}\cdots\text{Cg1}^{\text{i}}$	0.95	2.83	3.6180 (15)	141
$\text{C86}-\text{H86}\cdots\text{Cg1}^{\text{ii}}$	0.95	2.83	3.6614 (13)	147

Symmetry codes: (i)  $-x + \frac{3}{2}, -y + \frac{3}{2}, -z + 1$ ; (ii)  $x + \frac{3}{2}, y + \frac{3}{2}, z + 1$ .

Data collection: APEX2 (Bruker, 2004); cell refinement: APEX2 and SAINT (Bruker, 2004); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008) and OSCAIL (McArdle *et al.*, 2004); molecular graphics: PLATON (Spek, 2009); software used to prepare material for publication: SHELXL97.

CFL thanks the FCT and the European Social Fund (ESF) under the third Community Support Framework (CSF) for the award of a PhD Research Grant (SRFH/BD/29394/2006).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH5183).

## References

- Bruker (2004). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Lima, C. F., Gomes, L. R., Santos, L. M. N. B. F. & Low, J. N. (2010). *Acta Cryst. E* **66**, o3289.  
 McArdle, P., Gilligan, K., Cunningham, D., Dark, R. & Mahon, M. (2004). *CrystEngComm*, **6**, 303–309.  
 Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.  
 Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.

## supporting information

*Acta Cryst.* (2011). E67, o66 [https://doi.org/10.1107/S160053681005083X]

**4,4'-(1,8-Naphthalene-1,8-diyl)dibenzonitrile**

**Carlos F. Lima, Ligia R. Gomes, Luís M. N. B. F. Santos and John Nicolson Low**

**S1. Comment**


The exterior C1-C9-C8 angle of the naphthalene ring in C<sub>24</sub>H<sub>14</sub>N<sub>2</sub> is significantly larger, 126.106 (11)°, than that found in the two independent molecules of the single phenyl substituted compound, 4-(1-naphthyl)benzonitrile (Lima *et al.*, 2010), with values of 123.17 (11)° and 123.21 (10)° as are the angles C9—C1—C11 and C9—C8—C81, 124.93 (10)° and 124.79 (11)° as compared to the values for the single phenyl substituent of 121.463 (11)° and 121.47 (10)°.

The dihedral angles between the mean planes of the naphthalene ring and the C11—C16 ring and the C81—C86 rings are 66.33 (5)° and 67.89 (5)° respectively. These angles are significantly larger than those found for the single phenyl substituent in the two molecules of 4-(1-naphthyl)benzonitrile in which the naphthalene rings form dihedral angles of 60.28 (3)° and 60.79 (3)° for molecules 1 and 2 respectively.

C12 and C82 are linked *via* C—H $\cdots$  $\pi$  interactions to the centres-of-gravity of the rings C8—C10 and C1—C10 at (3/2 - x, 3/2 - y, 1 - y) respectively and C16 and C86 are linked *via* C—H $\cdots$  $\pi$  interactions to the centres-of-gravity of the rings C8—C10 and C1—C10 at (1 - x, 1 - y, 1 - y) respectively, Table 1. The molecules are thus linked into ladders with the molecules being stacked alternately head-to-tail as the rungs with the cyano groups and atoms C4 and C5 of the naphthalene groups pointing outwards. Alternate ladders run parallel to (110) and (-110). There is a solvent accessible void of 47 Å<sup>3</sup> in the structure lying between the ladders. This contains no residual electron density. There is no  $\pi\cdots\pi$  stacking nor are there C—H $\cdots$ N hydrogen bonds.


**S2. Refinement**

H atoms were treated as riding atoms with C—H(aromatic), 0.95 Å, with  $U_{\text{iso}} = 1.2U_{\text{eq}}(\text{C})$ . The positions of the H atoms were calculated and checked on a difference map during the refinement.



**Figure 1**

The molecular structure of the title compound with our numbering scheme. Displacement ellipsoids are drawn at the 30% probability level.



**Figure 2**

Stereoview of the ladders formed by C—H... $\pi$  interactions (dashed lines). Hydrogen atoms not involved in the motifs are not included.

## 4,4'-(1,8-Naphthalene-1,8-diyl)dibenzonitrile

## Crystal data

$C_{24}H_{14}N_2$	$F(000) = 1376$
$M_r = 330.37$	$D_x = 1.273 \text{ Mg m}^{-3}$
Monoclinic, $C2/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: $-C 2yc$	Cell parameters from 205 reflections
$a = 17.0872 (9) \text{ \AA}$	$\theta = 7.4\text{--}29.2^\circ$
$b = 8.2997 (4) \text{ \AA}$	$\mu = 0.08 \text{ mm}^{-1}$
$c = 24.3656 (13) \text{ \AA}$	$T = 150 \text{ K}$
$\beta = 93.795 (2)^\circ$	Plate, white
$V = 3447.9 (3) \text{ \AA}^3$	$0.40 \times 0.30 \times 0.02 \text{ mm}$
$Z = 8$	

## Data collection

Bruker SMART APEX diffractometer	11422 measured reflections
Radiation source: fine-focus sealed tube	4634 independent reflections
Graphite monochromator	3482 reflections with $I > 2\sigma(I)$
Detector resolution: $8.33 \text{ pixels mm}^{-1}$	$R_{\text{int}} = 0.031$
$\omega$ scans	$\theta_{\text{max}} = 29.2^\circ$ , $\theta_{\text{min}} = 3.0^\circ$
Absorption correction: multi-scan ( <i>SADABS</i> ; Bruker, 2004)	$h = -22 \rightarrow 23$
$T_{\text{min}} = 0.971$ , $T_{\text{max}} = 0.999$	$k = -11 \rightarrow 6$
	$l = -24 \rightarrow 33$

## Refinement

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.048$	H-atom parameters constrained
$wR(F^2) = 0.131$	$w = 1/[\sigma^2(F_o^2) + (0.0635P)^2 + 1.3141P]$
$S = 1.04$	where $P = (F_o^2 + 2F_c^2)/3$
4634 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
235 parameters	$\Delta\rho_{\text{max}} = 0.38 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.26 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	

## Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
N14	0.69235 (9)	0.27151 (19)	0.76246 (5)	0.0427 (4)
N84	0.60515 (8)	0.8449 (2)	0.77878 (5)	0.0449 (4)
C1	0.64595 (7)	0.48335 (15)	0.49000 (5)	0.0182 (3)
C2	0.66041 (7)	0.35609 (16)	0.45575 (5)	0.0216 (3)

H2	0.6753	0.2553	0.4717	0.026*
C3	0.65391 (8)	0.36994 (17)	0.39800 (5)	0.0237 (3)
H3	0.6644	0.2800	0.3755	0.028*
C4	0.63235 (8)	0.51393 (17)	0.37489 (5)	0.0228 (3)
H4	0.6277	0.5238	0.3360	0.027*
C5	0.59492 (7)	0.79704 (17)	0.38171 (5)	0.0225 (3)
H5	0.5909	0.8029	0.3427	0.027*
C6	0.57980 (8)	0.93033 (17)	0.41173 (5)	0.0238 (3)
H6	0.5655	1.0288	0.3939	0.029*
C7	0.58549 (7)	0.92080 (16)	0.46957 (5)	0.0219 (3)
H7	0.5755	1.0148	0.4902	0.026*
C8	0.60507 (7)	0.77995 (15)	0.49740 (5)	0.0183 (3)
C9	0.62271 (7)	0.63717 (15)	0.46689 (5)	0.0173 (3)
C10	0.61666 (7)	0.64945 (16)	0.40782 (5)	0.0192 (3)
C11	0.65558 (7)	0.44683 (15)	0.55012 (5)	0.0182 (3)
C12	0.71645 (7)	0.51380 (15)	0.58382 (5)	0.0199 (3)
H12	0.7517	0.5876	0.5687	0.024*
C13	0.72607 (7)	0.47384 (16)	0.63911 (5)	0.0220 (3)
H13	0.7670	0.5212	0.6620	0.026*
C14	0.67492 (8)	0.36320 (16)	0.66076 (5)	0.0226 (3)
C15	0.61429 (8)	0.29422 (16)	0.62760 (6)	0.0235 (3)
H15	0.5795	0.2194	0.6427	0.028*
C16	0.60519 (8)	0.33568 (16)	0.57243 (5)	0.0222 (3)
H16	0.5643	0.2880	0.5496	0.027*
C81	0.60545 (7)	0.78930 (15)	0.55880 (5)	0.0180 (3)
C82	0.66047 (8)	0.88695 (16)	0.58767 (5)	0.0213 (3)
H82	0.6978	0.9448	0.5682	0.026*
C83	0.66135 (8)	0.90069 (16)	0.64442 (5)	0.0227 (3)
H83	0.6997	0.9656	0.6638	0.027*
C84	0.60552 (8)	0.81858 (17)	0.67278 (5)	0.0227 (3)
C85	0.54901 (7)	0.72304 (17)	0.64435 (5)	0.0228 (3)
H85	0.5106	0.6680	0.6637	0.027*
C86	0.54928 (7)	0.70913 (16)	0.58781 (5)	0.0201 (3)
H86	0.5108	0.6443	0.5685	0.024*
C141	0.68499 (8)	0.31483 (19)	0.71759 (6)	0.0287 (3)
C841	0.60570 (8)	0.83269 (19)	0.73182 (6)	0.0297 (3)

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
N14	0.0462 (8)	0.0557 (10)	0.0262 (7)	-0.0037 (7)	0.0020 (6)	0.0074 (6)
N84	0.0411 (8)	0.0708 (11)	0.0227 (7)	-0.0121 (7)	0.0020 (6)	0.0000 (7)
C1	0.0156 (5)	0.0202 (6)	0.0184 (6)	-0.0023 (5)	-0.0006 (4)	-0.0009 (5)
C2	0.0210 (6)	0.0199 (7)	0.0239 (7)	0.0000 (5)	0.0007 (5)	-0.0011 (5)
C3	0.0229 (6)	0.0258 (7)	0.0227 (7)	-0.0014 (5)	0.0042 (5)	-0.0083 (5)
C4	0.0214 (6)	0.0298 (7)	0.0173 (6)	-0.0028 (5)	0.0018 (5)	-0.0033 (5)
C5	0.0210 (6)	0.0294 (7)	0.0169 (6)	-0.0014 (5)	0.0001 (5)	0.0039 (5)
C6	0.0216 (6)	0.0244 (7)	0.0253 (7)	0.0025 (5)	0.0020 (5)	0.0061 (5)

C7	0.0202 (6)	0.0210 (7)	0.0247 (7)	0.0007 (5)	0.0036 (5)	-0.0012 (5)
C8	0.0157 (5)	0.0221 (7)	0.0171 (6)	-0.0008 (5)	0.0023 (4)	-0.0016 (5)
C9	0.0142 (5)	0.0215 (6)	0.0161 (6)	-0.0009 (4)	0.0004 (4)	-0.0008 (5)
C10	0.0155 (5)	0.0248 (7)	0.0173 (6)	-0.0027 (5)	0.0013 (4)	-0.0001 (5)
C11	0.0188 (6)	0.0175 (6)	0.0182 (6)	0.0033 (5)	0.0014 (4)	-0.0010 (5)
C12	0.0182 (6)	0.0204 (6)	0.0209 (6)	0.0007 (5)	0.0006 (5)	0.0011 (5)
C13	0.0202 (6)	0.0240 (7)	0.0211 (6)	0.0016 (5)	-0.0025 (5)	-0.0017 (5)
C14	0.0256 (6)	0.0235 (7)	0.0188 (6)	0.0043 (5)	0.0021 (5)	0.0009 (5)
C15	0.0244 (6)	0.0213 (7)	0.0249 (7)	-0.0006 (5)	0.0033 (5)	0.0032 (5)
C16	0.0213 (6)	0.0207 (7)	0.0241 (7)	-0.0026 (5)	-0.0006 (5)	-0.0007 (5)
C81	0.0193 (6)	0.0170 (6)	0.0179 (6)	0.0034 (5)	0.0023 (4)	-0.0013 (5)
C82	0.0235 (6)	0.0202 (6)	0.0206 (6)	-0.0011 (5)	0.0033 (5)	0.0001 (5)
C83	0.0247 (6)	0.0213 (7)	0.0217 (6)	-0.0016 (5)	-0.0009 (5)	-0.0022 (5)
C84	0.0250 (6)	0.0253 (7)	0.0179 (6)	0.0026 (5)	0.0022 (5)	0.0008 (5)
C85	0.0202 (6)	0.0264 (7)	0.0222 (6)	0.0006 (5)	0.0041 (5)	0.0017 (5)
C86	0.0179 (6)	0.0214 (7)	0.0212 (6)	-0.0001 (5)	0.0020 (5)	-0.0025 (5)
C141	0.0290 (7)	0.0336 (8)	0.0235 (7)	-0.0007 (6)	0.0019 (5)	0.0020 (6)
C841	0.0277 (7)	0.0384 (9)	0.0229 (7)	-0.0045 (6)	0.0011 (5)	0.0005 (6)

*Geometric parameters (Å, °)*

N14—C141	1.1499 (19)	C11—C12	1.3971 (17)
N84—C841	1.1496 (19)	C12—C13	1.3867 (17)
C1—C2	1.3785 (18)	C12—H12	0.9500
C1—C9	1.4405 (17)	C13—C14	1.3951 (19)
C1—C11	1.4943 (17)	C13—H13	0.9500
C2—C3	1.4088 (18)	C14—C15	1.3938 (18)
C2—H2	0.9500	C14—C141	1.4411 (18)
C3—C4	1.3614 (19)	C15—C16	1.3864 (18)
C3—H3	0.9500	C15—H15	0.9500
C4—C10	1.4176 (18)	C16—H16	0.9500
C4—H4	0.9500	C81—C82	1.3957 (17)
C5—C6	1.3603 (19)	C81—C86	1.3976 (18)
C5—C10	1.4185 (18)	C82—C83	1.3864 (18)
C5—H5	0.9500	C82—H82	0.9500
C6—C7	1.4085 (18)	C83—C84	1.3926 (19)
C6—H6	0.9500	C83—H83	0.9500
C7—C8	1.3814 (18)	C84—C85	1.3973 (18)
C7—H7	0.9500	C84—C841	1.4431 (18)
C8—C9	1.4413 (17)	C85—C86	1.3829 (18)
C8—C81	1.4977 (17)	C85—H85	0.9500
C9—C10	1.4396 (17)	C86—H86	0.9500
C11—C16	1.3965 (18)		
C2—C1—C9	119.88 (11)	C13—C12—H12	119.6
C2—C1—C11	115.18 (11)	C11—C12—H12	119.6
C9—C1—C11	124.93 (11)	C12—C13—C14	119.19 (12)
C1—C2—C3	122.46 (12)	C12—C13—H13	120.4

C1—C2—H2	118.8	C14—C13—H13	120.4
C3—C2—H2	118.8	C15—C14—C13	120.76 (12)
C4—C3—C2	119.08 (12)	C15—C14—C141	118.65 (13)
C4—C3—H3	120.5	C13—C14—C141	120.58 (12)
C2—C3—H3	120.5	C16—C15—C14	119.44 (12)
C3—C4—C10	121.23 (12)	C16—C15—H15	120.3
C3—C4—H4	119.4	C14—C15—H15	120.3
C10—C4—H4	119.4	C15—C16—C11	120.60 (12)
C6—C5—C10	120.96 (12)	C15—C16—H16	119.7
C6—C5—H5	119.5	C11—C16—H16	119.7
C10—C5—H5	119.5	C82—C81—C86	118.93 (11)
C5—C6—C7	119.30 (12)	C82—C81—C8	119.46 (11)
C5—C6—H6	120.4	C86—C81—C8	121.52 (11)
C7—C6—H6	120.4	C83—C82—C81	120.86 (12)
C8—C7—C6	122.49 (12)	C83—C82—H82	119.6
C8—C7—H7	118.8	C81—C82—H82	119.6
C6—C7—H7	118.8	C82—C83—C84	119.47 (12)
C7—C8—C9	119.65 (11)	C82—C83—H83	120.3
C7—C8—C81	115.56 (11)	C84—C83—H83	120.3
C9—C8—C81	124.79 (11)	C83—C84—C85	120.37 (12)
C10—C9—C1	116.95 (11)	C83—C84—C841	119.90 (12)
C10—C9—C8	116.99 (11)	C85—C84—C841	119.73 (12)
C1—C9—C8	126.06 (11)	C86—C85—C84	119.56 (12)
C4—C10—C5	119.01 (12)	C86—C85—H85	120.2
C4—C10—C9	120.40 (12)	C84—C85—H85	120.2
C5—C10—C9	120.59 (12)	C85—C86—C81	120.79 (12)
C16—C11—C12	119.22 (12)	C85—C86—H86	119.6
C16—C11—C1	119.01 (11)	C81—C86—H86	119.6
C12—C11—C1	121.67 (11)	N14—C141—C14	177.90 (17)
C13—C12—C11	120.78 (12)	N84—C841—C84	179.27 (18)
C9—C1—C2—C3	0.44 (19)	C2—C1—C11—C12	111.39 (14)
C11—C1—C2—C3	179.92 (11)	C9—C1—C11—C12	-69.16 (16)
C1—C2—C3—C4	-0.18 (19)	C16—C11—C12—C13	-1.45 (19)
C2—C3—C4—C10	0.18 (19)	C1—C11—C12—C13	-177.65 (12)
C10—C5—C6—C7	-0.33 (19)	C11—C12—C13—C14	1.13 (19)
C5—C6—C7—C8	-0.80 (19)	C12—C13—C14—C15	-0.6 (2)
C6—C7—C8—C9	1.79 (19)	C12—C13—C14—C141	177.82 (13)
C6—C7—C8—C81	-177.71 (11)	C13—C14—C15—C16	0.4 (2)
C2—C1—C9—C10	-0.67 (17)	C141—C14—C15—C16	-178.07 (13)
C11—C1—C9—C10	179.91 (11)	C14—C15—C16—C11	-0.7 (2)
C2—C1—C9—C8	179.44 (12)	C12—C11—C16—C15	1.22 (19)
C11—C1—C9—C8	0.02 (19)	C1—C11—C16—C15	177.52 (12)
C7—C8—C9—C10	-1.62 (17)	C7—C8—C81—C82	-65.72 (15)
C81—C8—C9—C10	177.84 (11)	C9—C8—C81—C82	114.80 (14)
C7—C8—C9—C1	178.27 (11)	C7—C8—C81—C86	110.83 (14)
C81—C8—C9—C1	-2.27 (19)	C9—C8—C81—C86	-68.65 (16)
C3—C4—C10—C5	179.57 (12)	C86—C81—C82—C83	2.07 (19)

C3—C4—C10—C9	-0.45 (19)	C8—C81—C82—C83	178.71 (12)
C6—C5—C10—C4	-179.61 (12)	C81—C82—C83—C84	-1.37 (19)
C6—C5—C10—C9	0.41 (19)	C82—C83—C84—C85	0.0 (2)
C1—C9—C10—C4	0.68 (16)	C82—C83—C84—C841	-179.70 (13)
C8—C9—C10—C4	-179.42 (11)	C83—C84—C85—C86	0.64 (19)
C1—C9—C10—C5	-179.34 (11)	C841—C84—C85—C86	-179.66 (12)
C8—C9—C10—C5	0.56 (17)	C84—C85—C86—C81	0.08 (19)
C2—C1—C11—C16	-64.81 (15)	C82—C81—C86—C85	-1.42 (18)
C9—C1—C11—C16	114.64 (14)	C8—C81—C86—C85	-177.98 (11)

*Hydrogen-bond geometry (Å, °)*

Cg1 and Cg2 are the centroids of the C1—C10 and C8—C10 rings, respectively.

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
C12—H12 $\cdots$ Cg2 <sup>i</sup>	0.95	2.75	3.6147 (13)	152
C16—H16 $\cdots$ Cg2 <sup>ii</sup>	0.95	2.92	3.6539 (15)	135
C82—H82 $\cdots$ Cg1 <sup>i</sup>	0.95	2.83	3.6180 (15)	141
C86—H86 $\cdots$ Cg1 <sup>ii</sup>	0.95	2.83	3.6614 (13)	147

Symmetry codes: (i)  $-x+3/2, -y+3/2, -z+1$ ; (ii)  $x+3/2, y+3/2, z+1$ .