

# Oral Microbiome Associated with Peri-Implant Infections – A Review

Molina F.<sup>1</sup>, Lopes Cardoso I.<sup>2</sup>, Moura Teles A.<sup>3</sup>, Pina C.<sup>4</sup>

<sup>1</sup>Student of the Integrated Master in Dental Medicine, Health Sciences Faculty, Fernando Pessoa University, Rua Carlos da Maia, 296, 4200-150 Porto, Portugal

<sup>2,3,4</sup>Health Sciences Faculty, Fernando Pessoa University, Rua Carlos da Maia, 296, 4200-150 Porto, Portugal

**Abstract:** *Dental treatments using dental implants have been well documented over the past 40 years and with great success. The dental implant installed in the place of missing teeth should always involve proper forecasting by the dentist. Namely, it is important to know the microbiome surrounding the implant, from its planning till final rehabilitation. The exact time of microbiome formation, as well as microorganisms involved, are essential for the proper implementation and success of the implant. However, internal contaminations of the rehabilitated implants, the extracellular components of microorganisms, such as endotoxins, have a huge influence on implant success. In addition, it is also very important the knowledge concerning implants surfaces and associated microorganisms. This study conducted a literature review on the oral microbiome and its relationship with the peri-implant infection, with the discussion of several classical and current studies. Although it can be concluded that the peri-implant microbiome is characterized by the microbiome present before dental implant placing, more studies are required to better elucidate the planning and the longevity of dental implant treatment.*

**Keywords:** dental implant; peri-implant microbiome; oral microbiome; periodontal disease

## 1. Introduction

Dental implants have been largely used for the last 40 years. They are artificial structures, made up of titanium, that are installed in the place of a missing tooth. Implants are anchored in the bone, through the concept of osseointegration [1].

The implant-abutment connection represents the weakest point of the dental implant, due to a micro-gap between the implant-abutment interface, which may cause microbial leakage [2]. The occurrence of microleakage through this interface is not surprising since the diameter of the smallest oral bacterial species may range from 0.2 to 0.7  $\mu\text{m}$  [3] and the micro-gaps resulting from implant-abutment attachment vary from 1 to 70  $\mu\text{m}$  [4], depending on the type of platform connection [5]. These hollow spaces may act as reservoir for commensal and/or pathogenic bacteria, especially anaerobic or micro-aerophilic species, representing a potential source of tissue inflammation, which may lead to bone resorption [6]. In consequence of that bacterial leakage, two types of diseases, peri-implant mucositis and peri-implantitis, that, being like gingivitis and periodontitis, can affect implants.

Mucositis is characterized by inflammation restricted to the peri-implant mucosa, without bone loss, while peri-implantitis is a deep inflammatory lesion, characterized by bleeding, peri-implant pocket and progressive loss of bone support around implants [7,8]. Peri-implantitis, like periodontitis, is an endogenous infection, poly-microbial and opportunist, that occurs by the conjunction action of local microbiota [7].

This study involves a bibliographic review concerning oral microbiome formation that occurs around implants and its characterization, as well as the analysis of peri-implant infections that can evolve to the undesirable implant loss.

The theme is of great importance in nowadays rehabilitation scenery since it is commonly performed in dental practices and is well accepted by patients. Therefore, clinicians, in general, must possess knowledge to perform an adequate and safe procedure and ensure long lasting dental implants treatments.

## 2. Materials and Methods

A bibliographic search was performed using *PubMed* and *B-On* databases. Keywords used were: “dental implants AND microbiology”, “loss of dental implants AND microbiology”, “microbiology AND microleakage”, “oral microbiome”.

### Oral Microbiome

The oral cavity is a dynamic system continuously colonized by microorganisms being called the oral microbiome. This recent concept assumes the involvement of these microorganisms with the host, including the survival capacity against the immune system defence [9].

In the oral cavity, the microbiome is inserted in different anatomic structures, in the form of biofilm, forming an ecosystem that, in equilibrium, allows the maintenance of health. However, ecological changes in the microbiome allow the development of pathogenic agents that can cause disease [10].

Biofilm development occurs in teeth and mucosa, as well as in artificial surfaces like prosthesis and implants. Therefore, oral biofilm acts as a pathogenic community, where microorganisms become less vulnerable to immune cells, like neutrophils and antibodies as well as anti-microbial agents [11]. Nowadays, the idea of how oral biofilm causes disease, like caries, periodontitis or peri-implantitis disease is explained by the ecological hypothesis of plaque, that

determines that it is the interaction between microorganisms and the host that defines the state of health or disease [12].

Recent advances from metagenomic studies have developed a new model of periodontal disease pathogenesis. Chronic periodontitis does not result from individual pathogens but rather from polymicrobial synergy and dysbiosis associated with a dysregulated immune response inducing inflammation-mediated tissue damage [13]. Namely, changes in microflora environment, due to pH, diet and hygiene alterations, the use of broad range antibiotics, the commitment of the immune system, among others, can lead to changes in biofilm composition. Because of those changes, certain microbial species can exhibit higher virulence, allowing opportunist microorganisms to cause disease [12,13,14].

The way tissues are formed around implants, when compared with teeth, is different, however similarities exist in the microbial colonization during biofilm formation [15]. Despite this similarity, fundamental differences exist between periodontal and peri-implant tissues, like absence of Sharpey fibres in peri-implant tissues [9]. In this case, collagen fibres of the submucous connective tissue are aligned in parallel to implant surfaces, resulting in a deeper gap than the gingival sulcus, allowing easier penetrance of microorganisms.

The lack of periodontal ligament gives certain disadvantages to the implant when compared with teeth, like:

- Reduction of physical barriers that allows entrance of microorganisms in submucous tissues; consequently, the conformation of peri-implant tissues makes them more susceptible to endogenous infections, when compared with periodontal tissues;
- Lack of adequate blood flow that decreases supply of nutrients and immune system cells, required for early stages of infection;
- Reduction of implants response to masticatory forces [8].

In face of this scenario, immediate postoperative failures can occur, resulting from contamination during surgery, that show up as microbial infections and healing problems [9]. Late failure usually occurs as a result of chronic infections in peri-implant tissues. It is therefore of great importance the study of the oral microbiome that is present in dental implants and the way it sets up.

### 3. Development of peri-implant microbiome

The use of osseointegration has shown to be an excellent method in the replacement of lost teeth [16-21]. Osseointegrated dental implants (Figure 1) are metal structures, mainly made up of titanium, that are surgically installed in the alveolar bone, in the place where one or more teeth were lost. They are used due to their characteristics like biocompatibility, stability and resistance to corrosion.




Figure 1: Recently installed implant.

After dental implant installation, and according with the circumstances and planning of the case, a prosthetic restoration is installed, with the help of prosthetic components that are linked to implants and will support the prosthesis, giving the required function and aesthetics [9].

Prosthetic components (Figure 2) are mostly made up of metal, however, and for aesthetic reasons, other materials have been introduced in the prosthetic components of dental rehabilitations like zirconia oxide [22]. Differences in microbial colonization between ceramic and metal surfaces are still a controversial issue, as some *in vitro* studies point out that microbial colonization in zirconia oxide surfaces is lower when compared with titanium surfaces [23,24] and others verified that the predominant factor for adhesion depends more on the bacterial species itself [25].



Figure 2: Examples of prosthetic abutments.

With the increase in the use of dental implants rehabilitation, it becomes necessary the study on biofilm formation, composition and action on implant surface. The formation of dental biofilm on implants surface can lead to the development of local infections capable of causing implant loss [9]. In this way, it is very important the study of dental biofilm formation on implants and understand when it occurs.

Several studies demonstrate biofilm formation on dental implants. Koka et al. [26], in a study with partially edentulous patients, suggested that colonization of marginal implant plaque occurred within 14 days, whereas subgingival colonization took longer and occurred within 28 days (Figure 3) [26].



**Figure 3:** Second surgical step, 11<sup>th</sup> region, installed healer.

The period of peri-implant microbiome establishment has been widely discussed and there is still no consensus concerning the exact time of its start. Different methodologies for species identification, probably can justify several results.

In 1996, Persson *et al.* [27] observed that implant contamination occurred during the first and second surgical steps (installation of prosthetic components) which agrees with data from van Winkelhoff & Winkel [28] who concluded that microbial flora is present after implant installation, with the presence of *Peptostreptococcus micros*, *Fusobacterium sp.* and *Prevotella intermedia*.

Bacteria associated with periodontitis (*Porphyromonas gingivalis*, *Tannerella forsythia* and *Treponema denticola*) were detected with the same prevalence 30 minutes or one year after implant installation, being *P. gingivalis* the most prevalent microorganism in teeth and implants [29].

Other studies indicate that, a few weeks after dental implants installation, it is possible to detect the presence of microorganisms in samples collected from the peri-implant groove or from implants [19,30,31].

Several authors suggest that the establishment of peri-implant microbiome occurs by transmission of remaining microbiome from teeth to implants [19,31,32]. The tongue can also be a source of microorganisms for implant colonization [33]. According with Gerber *et al.* [34], the oral microbiome present before implant installation can determine the composition of the microbiome in the peri-implant area. Patients with history of periodontal disease can have significant impact on peri-implant microbiome [34]. This might result in pathogen transmission from remaining teeth to implants [35].

In the meantime, patients with history of aggressive properly controlled periodontitis, that were rehabilitated with dental implants, did not have changes or inflammations after osseointegration [36]. On the other hand, Botero *et al.* [17] verified that partially edentulous patients with periodontal involvement, that were rehabilitated with dental implants, showed around implants, pocket and bone loss, with high levels of periodontopathogens (facultative anaerobes after 6 months of implant exposition to oral environment) in subgingival microbiome. This indicates that teeth might serve as a microbial reservoir, suggesting that patients with history of periodontal disease would have increased risk of peri-implant disease [17].

In this way, teeth and implants can serve as niches for dental biofilm formation and induce gingivitis and mucositis, that can develop to periodontitis and peri-implantitis respectively [19].

#### 4. Dental implant surfaces

Artificial surfaces of implants and rehabilitation structures allow the formation of dental biofilm [9]. Therefore, it is required to know surfaces of different types of implants for an adequate choice for lowest possible biofilm accumulation.

Concerning implant surface, there is no consensus among scientific community on if microbial composition is affected by different geometry implants' surfaces. Some studies report relevant results but are constantly changing due to new technologies in dental implants manufacture. Several changes in implant design have been done in the last years, in an attempt to decrease the space between the implant and the prosthetic component, to reduce bacterial proliferation. However, only limited success was obtained so far [37].

Rougher implants can favour biofilm development and consequently peri-implantitis [38]. In a more recent study it was also shown that peri-implantitis can lead to bigger bone loss, due to the presence of periodontal pathogens in implants of internal connexion, when compared to implants with "morse cone" type connexions [37].

Besides different implant surfaces that suffer microbial colonization, implants are also subjected to local exudates, pH, toxins and components of microbial release. In this way, some authors study titanium corrosion and its consequences, namely, the tribocorrosion which is a material degradation process due to the combined effect of corrosion and wear and that has been identified as the major degradation mechanism that results in dental implants failure [39]. Biological consequences of titanium corrosion and its accumulation in tissues are significant, and there may be changes in peri-implant tissues and chronic inflammation that can lead to implant loss [40]. Titanium corrosion has been detected at considerable levels in tissues and fluids adjacent to implants [41].

In fact, physiology of the environment where the implant is inserted varies with the place and its installation mode, being composed by a complex system with organic and inorganic

components, like ions, aminoacids, proteins and fatty acids, as well as living cells [42].

The composition of that environment can be altered according with patient's health and, consequently, can lead to implant loss [43]. Therefore, inflammation and/or conditions associated with corrosion can create an acidic environment [44]. Measurements done in titanium alloys of orthopaedic implants showed that implant's place is acid, with pH around 2,5. For dental implants, this can be associated to the presence of biofilm or acidogenic bacteria, capable of reducing local pH below 4,5 [45].

So, debris released from the degradation of dental implants has cytotoxic and genotoxic potential for peri-implant tissues. Thus, the amount and physicochemical properties of the degradation products determine the magnitude of the detrimental effect on peri-implant tissues [46].

## 5. Oral microbiome associated with dental implants

Subgingival microbiome associated with dental implants has shown to be similar to dental microbiome [47,48]. However, microbiological studies have suggested the transmission of subgingival microorganisms, from dental locations to peri-implant spots [49].

Initial colonization of the oral microbiome occurs by adhesion of earlier colonizers (earlier microorganisms), including *Streptococcus oralis*, *S. gordonii* and *Actinomyces naeslundii*. This initial colonization favours adhesion of secondary invaders where biofilm formation occurs by the interaction and multiplication of available microorganisms [50]. With biofilm establishment, synergic and antagonist microbial interactions occur enriching the biofilm in the spot and in prosthetic components, making clear the relationship between the biofilm and host response [50].

Biofilm in healthy peri-implant spots has been characterized with high proportion of Gram positive coccus. In the meantime, in peri-implantitis cases, biofilm has high amounts of Gram negative bacteria, as well as Gram positive coccus (*Parvimonas* sp. and *Peptostreptococcus* sp.) [51]. However, other studies reported that *Aggregatibacter actinomycetemcomitans* and *P. gingivalis* were found in big quantity in peri-implant lesions [52,53]. These two pathogens can be considered the predominant microorganisms, being responsible for destructive infection in peri-implantitis [17]. Also, Van de Velde et al. [54] demonstrated the presence of *A. actinomycetemcomitans*, *Fusobacterium* sp., *P. gingivalis*, *Pseudomonas aeruginosa* and *T. forsythia*, in implants diagnosed with peri-implantitis. Symbiosis between *Bacteroides* sp. and *P. aeruginosa* seems to favour the persistence of *P. aeruginosa* in inflamed regions around implants [54].

Researchers did not observe considerable differences in subgingival flora around teeth or dental implants [30]. Other studies also analysed the frequency of *A. actinomycetemcomitans* and *A. viscosus* in supragingival biofilm, observing 92% vs 57% in teeth and 90% vs 73% in

implants, suggesting that subgingival microbiome is similar in teeth and implants [55]. In 2006, Quirynen et al. [19], using molecular biology techniques like PCR, observed small differences in microflora of teeth and implants [19].

In partially edentulous patients, morphological types found did not show significative differences between teeth and implants. Kohavi et al. [55] identified similar microbiome in healthy teeth of partially and totally edentulous patients, being mainly composed of Gram-positive coccus with low count of spirochetes and mobile bacilli.

However, in 2011, Quirynen & Van Assche [56] detected high levels of bacteria related with periodontitis and peri-implantitis, in totally edentulous patients, agreeing with other similar studies [30,57,58].

Periodontal bacteria, like *P. gingivalis*, *T. forsythia* and *T. denticola*, were found with the same prevalence, in teeth and implants, being *P. gingivalis* the most prevalent [29]. It was also observed high prevalence of *Fusobacterium* sp. in implants spots after their installation, agreeing with data from van Winkelhoff & Winkel [28].

In 2008, Shibli et al. [59] evaluated the supra and subgingival microflora, in healthy implants and implants with peri-implant disease. Supragingival microflora was similar in both groups [59]. *Veilonella parvula* and *F. periodonticum* were the dominant species [59]. Besides these, four bacterial species were found in high significance levels in the group of implants with peri-implantitis: 3 periodontal pathogens of the red complex (*P. gingivalis*, *T. forsythia* and *T. denticola*) and *P. nigrescens* ( $P < 0,05$ ) [59].

In subgingival region, the complex profiles that shelter most beneficial species (purple, yellow and green) were similar between healthy implants and implants with peri-implantitis. Most pathogens of the red and orange complexes were found in high levels in the group of implants having peri-implantitis [60,61]. *P. gingivalis*, *T. forsythia*, *T. denticola*, *F. nucleatum*, *F. nucleatum ss vicentii* and *P. intermedia* were found in significantly higher levels in subgingival biofilm of implants with peri-implantitis ( $p < 0,05$ ) [60,61]. Three bacterial species (*A. naeslundii*, *S. intermedius* and *S. mitis*) and a facultative periodontal pathogen (*F. periodonticum*) were found in high levels in supragingival biofilm when compared with subgingival samples of healthy implants ( $p < 0,05$ ) [59].

Levels of 3 microorganisms, *V. parvula*, *S. gordonii* and *S. intermedius*, as well as *F. periodonticum*, were significantly higher in the supragingival biofilm when compared with subgingival biofilm of implants with peri-implantitis. A higher medium count was observed for certain pathogens like *F. nucleatum nucleatum*, *P. intermedia*, *P. nigrescens*, *T. denticola*, *Selenomonas noxia* ( $p > 0,05$ ) and *T. forsythia* ( $p < 0,05$ ) in subgingival biofilm of implants with peri-implantitis [59].

In 1991, Slots et al. [62] also reported that in implants with peri-implantitis it is possible to detect big quantities of Gram-negative anaerobic bacteria, including Fusobacteria, spirochetes, *B. forsythus*, *P. intermedia*, *P. nigrescens* and *P.*

*gingivalis*. *A. actinomycetemcomitans* was also isolated in this type of lesion. In this way, microflora in peri-implantitis lesions resemble adult refractory periodontitis [62].

Recently, Zhuang *et al.* [63] demonstrated that periodontal pathogens are common in both places, periodontal and peri-implant, independently of health or disease. It was also evident that there was difference in the involvement of some pathogens in disease conditions. Prevalence and level of *P. gingivalis* and *F. nucleatum* were significantly associated with periodontitis, but not associated with peri-implantitis. However, *A. actinomycetemcomitans* was associated with periodontitis as well as with peri-implantitis [63].

It is important to note that differences observed in these studies can be explained by the use of different methodologies in microbial identification, since half of the oral biofilm is composed of non-cultivable microflora [64].

As a conclusion, differences seem to exist between microbial profile of healthy implants and implants with peri-implant disease, in supra and subgingival biofilm. The main differences were high levels of certain periodontal pathogens and low proportions of microbial complexes (compatible with the host) in the group of implants with peri-implantitis. The striking presence of species of the red complex in supragingival biofilm of implants with peri-implantitis suggests a propitious environment for reservoir of pathogenic species, being able to contribute to re-infection in treated subgingival spots [59].

## 6. Association of peri-implant disease with periodontal disease

Studies have shown that the microbiome around implants that failed is similar to the microbiome of teeth with periodontitis, in composition as well as proportion of microorganisms. Hence, the association of peri-implant disease with periodontal disease have been reported for several studies [37,49].

The incidence of peri-implantitis has increased continuously over the last years, being related with the loss of implants. Many factors play a fundamental role in complications and implant failures like implant and prosthetic components design [37] as well as bacterial colonization in implants.

Bacterial colonization around dental implants can lead to some local diseases, being the most common, mucositis and peri-implantitis. Mucositis (Figure 4) is an inflammation restricted to peri-implant mucosa in the implant, without bone loss. Peri-implantitis (Figure 5) is a deep inflammatory lesion, characterized by bleeding, peri-implant pocket and progressive loss of bone support around implants. These two forms of infection resemble respectively gingivitis and periodontitis in teeth [7,8].



Figure 4: Mucositis spots.



Figure 5: Periapical X-ray of implant affected by peri-implantitis.

Peri-implantitis, like periodontitis, is an endogenous polymicrobial and opportunist infection, that appears by the combined action of local microbiome [7], resulting from the disequilibrium between bacteria and host response, that can lead to local inflammation or bone loss, and sometimes to implant loss [8].

Several studies that indicate dental implant failure after osseointegration, show the presence of high levels of periodontopathogenic bacteria (*P. gingivalis*, *P. intermedia* and *A. actinomycetemcomitans*) in peri-implant lesions [17,26,65,66]. In patients with a history of periodontitis, putative periodontal pathogens prevailed in the microbiome of diseased implants. Diseased implants and corresponding healthy sites appear to have distinct microbiological ecosystems [67].

Bacterial colonization by periodontopathogens has been considered a risk factor for peri-implantitis. Other studies indicate that teeth can be a source of bacteria in partially edentulous patients that were rehabilitated with dental implants [35,68,69].

In the meantime, the presence of microorganisms related with periodontitis and peri-implantitis found in gingival and peri-implant grooves, respectively, did not mean implant failure, and did not result in disease [31].

Work of Renvert & Persson [49] showed that patients with periodontitis history can have increased risk of peri-implant

infections. Moreover, Heitz-Mayfield & Lang [8] stated that patients susceptible to periodontal disease, have higher susceptibility to peri-implant disease when compared with patients without history of periodontal disease [8].

To decrease the chance of implant loss due to periodontopathogenic bacteria, several researchers propose elimination of these pathogens before dental implant installation [28,70,71].

In individuals with history of periodontal disease, local decontamination prior to implant setting should be done, but also in patients without history of periodontal disease. Besides, scientific data are limited in relation to periodontopathogen colonization in patients without history of periodontal disease and little is known about the efficacy of reduction of these microorganisms, especially in long term [72,73]. Nowadays, it is believed that treated sites are subject to recolonization with a microbiota similar to that present before therapy. The degree and speed of recolonization depends on the treatment protocol, the distribution patterns of periodontal microorganisms elsewhere in the oral cavity and the quality of the patient's oral hygiene [74].

Plus, in classic bacterial infections, the diversity of the microbiota decreases as the disease develops. However, in most cases of periodontitis, the diversity of the flora increases. Given the large diversity and the complex interactions among the members of the microbiota, a therapeutic concept that targets one responsible bacterial species or strain with a highly specific agent appears to be an unrealistic approach. Therefore, little evidence supports microbiological testing as an approach to obtain better clinical outcomes. In fact, at present, no protocol exists with proven superiority, in terms of efficiency or effectiveness over scaling and root planning plus systemic amoxicillin and metronidazole for the therapy of any form of periodontal disease [74].

## 7. Dental Implant Connexions and Internal Contamination

Infiltration of microorganisms and their sub-products in implants makes relevant the study of implant and its prosthetic components surfaces. Penetration of bacteria and bacterial fluids can occur to the internal portion of dental implants and can be the cause of the inflammatory process that occurs in peri-implant tissues [68,75,76].

One of the major changes in the use of two steps implants has been the use of implants type "morse cone", to try avoiding internal contamination of implants, through the decrease of spaces between the implant and corresponding prosthetic components. However, microorganisms can grow and serve as bacterial stores in those places, being able to induce inflamed areas and bone loss around implants, through prosthetic components (in prosthetic pillar/implant junction) [27].

Therefore, implants are not protected against bacterial installation, becoming necessary changes in the connexion area of prosthetic pillar with the implant [77]. In a recent

meta-analysis study, Tallarico *et al.* [78] concluded that bacteria could easily be colonized at the implant-abutment interface and also affirmed that it is evident from a clinical point of view that inner portions of implant-abutment interfaces should always be considered contaminated, even in clinically healthy conditions.

Existing spaces between the retention screw and the prosthetic pillar are the main pathway of bacterial access in crowns on implants [79]. Through those spaces it is possible that fluids with bacterial sub-products and nutrients required for bacterial growth slip away by the interface prosthetic pillar/implant, contributing for the bad breath and the development of peri-implantitis [34].

Dental implant design in the connexion area of the implant with prosthetic pillar, might have strong impact on bacterial penetration [80].

Quirynen & van Steenberghe [81] showed bacterial penetration in the interface prosthetic pillar/implant, with implants of the external hexagon type.

Another work studied bacterial penetration through the interface prosthetic pillar/implant, in 13 different combinations of prosthetic pillar/implant [77]. Among diverse combinations used, the internal connexion and the use of silicone rings demonstrated to have lower bacterial infiltration [77].

The existence of spaces smaller than 4  $\mu\text{m}$  between crowns and implants of the *Ha-Ti* system is not an effective barrier against infiltration by *S. aureus* [82]. Work of Orsini *et al.* [85] was in accordance with that, mentioning spaces of 1 to 5  $\mu\text{m}$  between implant and the retention screw that could be filled by bacteria. Therefore, bacteria can be found in the most apical portion of the implant cylinder.

According with Piattelli *et al.* [84], prosthesis retained by screws or by cement can suffer penetration by bacteria or bacterial fluids to the inside of implants. However, there was less penetration in prosthesis retained by cement, being these ones more effective against penetration of bacteria and bacterial fluids [84].

Some implant systems possess sealing by a silicone ring that might reduce bacterial infiltration [85]. In this study, contamination of internal portions of prosthetic pillars of implants retained by screws was analysed. These researchers observed an amorphous and crystalline contamination suggesting calcium and phosphate, in all surfaces [85].

Microbial contamination was seen with higher intensity in the group without sealing [85]. No differences in bacterial morphology was observed between the group of sealed implants and the group without sealing. Most abundant microorganisms were coccus while bacilli were very rarely found. In clinical cases, microbial infiltration occurred in the interface prosthetic pillar/implant, although this contamination had been limited to patients having high standard of oral hygiene. In this way, authors concluded that contamination can be reduced with the use of silicone ring [85].

In the *in vitro* study of Steinebrunner *et al.* [80], bacterial infiltration was analysed along the interface prosthetic pillar/implant. Implants with internal connexions of the type tri-channels showed the higher amount of bacterial infiltration relative to implants of external hexagon connexion, internal hexagon and implants with silicone ring and implants with internal connexion by friction, respectively [80].

In other *in vitro* studies, without charges and charge dynamics, it was demonstrated that implants with “morse cone” connexion had minimal infiltration. These studies looked for contamination by *A. actinomycetemcomitans*, *P. gingivalis* and *E. coli* [86,87].

Microorganism penetration occurs in the different types of existing connexions. For this, it is required the development of new technologies for the improvement and decrease of internal contamination of implants [88].

## 8. Conclusion

From this review performed on oral microbiome and its relationship with peri-implant infections, it is possible to conclude that:

- Peri-implant microbiome is composed of microbiome present before installation of dental implants;
- The moment when microbiome develops on dental implant is still under discussion and there is no consensus among researchers;
- Microbiome established around dental implants is similar to the microbiome of periodontitis, in health, and also in cases of periodontal disease;
- There are several prosthetic components for rehabilitation of masticatory and aesthetical functions of the patient, however no totally efficient component is still available against micro-infiltrations to the inside of the implant;
- Deeper studies are still required to find an implant with the correct surface that decreases microbial colonization and ensures bigger success in dental implant treatments.

## References

- [1] P.I. Branemark, Osseointegration and its experimental background. *J. Prosthet. Dent.* 50(3) (1983) 399-410.
- [2] F. Mencia, F. de Angelis, P. Papi, D. Rosella, G. Pompa, S. di Carlo, A randomized clinical trial about presence of pathogenic microflora and risk of peri-implantitis: comparison of two different types of implant-abutment connections. *Eur. Rev. Med. Pharmacol. Sci.* 21 (2017) 1443-1451.
- [3] R.E. Barbosa, C. do Nascimento, J.P. Issa, E. Watanabe, I.Y. Ito, R.F.Jr. de Albuquerque, Bacterial culture and DNA checkerboard for the detection of internal contamination in dental implants. *J. Prosthodont.* 18(5) (2009) 376-381.
- [4] S. Harder, B. Dimarczek, Y. Açil, H. Terheyden, S. Freitag-Wolf, M. Kern, Molecular leakage at implant-abutment connection - in vitro investigation of tightness of internal conical implant-abutment connections against endotoxin penetration. *Clin. Oral Investig.* 14(4) (2010) 427-432.
- [5] A.R. de Freitas, T.S.O. Silva, R.F. Ribeiro, R.F. de Albuquerque Junior, V. Pedrazzi, C. do Nascimento, Oral bacterial colonization on dental implants restored with titanium or zirconia abutments: 6-month follow-up. *Clin. Oral Investig.* (2018) <https://doi.org/10.1007/s00784-018-2334-0>.
- [6] F. Mencia, P. Papi, S. di Carlo, G. Pompa, Salivary bacterial leakage into implant-abutment connections: preliminary results of an in vivo study. *Eur. Rev. Med. Pharmacol. Sci.* 20 (2016) 2476-2483.
- [7] G. Charalampakis, G.N. Belibasakis, Microbiome of peri-implant infections: lessons from conventional, molecular and metagenomic analyses. *Virulence.* 6(3) (2015) 183-187.
- [8] L.J. Heitz-Mayfield, N.P. Lang, Comparative biology of chronic and aggressive periodontitis vs. peri-implantitis. *Periodontol.* 2000. 53 (2010) 167-181.
- [9] G.N. Belibasakis, Microbiological and immunopathological aspects of peri-implant diseases. *Arch. Oral Biol.* 59(1) (2014) 66-72.
- [10] M.F. Zarco, T.J. Vess, G.S. Ginsburg, The oral microbiome in health and disease and the potential impact on personalized dental medicine. *Oral Dis.* 18(2) (2012) 109-120.
- [11] C. Schaudinn, A. Gorur, D. Keller, P.P. Sedghizadeh, J.W. Costerton, Periodontitis: an archetypical biofilm disease. *J. Am. Dent. Assoc.* 140(8) (2009) 978-986.
- [12] P.D. Marsh, Are dental diseases examples of ecological catastrophes? *Microbiology.* 149(Pt 2) (2003) 279-294.
- [13] Z.-L. Deng, S.P. Szafranski, M. Jarek, S. Bhujju, I. Wagner-Döbler, Dysbiosis in chronic periodontitis: key microbial players and interactions with the human host. *Sci. Rep.* 7(1) (2017) 3703.
- [14] V. Meuric, S. Le Gall-David, E. Boyer, L. Acuña-Amador, B. Martin, S.B. Fong, F. Barloy-Hubler, M. Bonnaure-Mallet, Signature of microbial dysbiosis in periodontitis. *Appl. Environ. Microbiol.* 83(14) (2017) e00462-17.
- [15] M.A. Listgarten, Clinical trials of endosseous implants: issues in analysis and interpretation. *Ann. Periodontol.* 2(1) (1997) 299-313.
- [16] R. Adell, B. Eriksson, U. Lekholm, P.I. Branemark, T. Jemt, Long-term follow-up study of osseointegrated implants in the treatment of totally edentulous jaws. *Int. J. Oral Maxillofacial Implants.* 5(4) (1990) 347-359.
- [17] J.E. Botero, A.M. González, R.A. Mercado, G. Olave, A. Contreras, Subgingival microbiota in peri-implant mucosa lesions and adjacent teeth in partially edentulous patients. *J. Periodontol.* 76(9) (2005) 1490-1495.
- [18] M. Bischof, R. Nedir, S. Abi Najm, S. Szmukler-Moncler, J. Samson, A five-year life-table analysis on wide neck ITI implants with prosthetic evaluation and radiographic analysis: results from a private practice. *Clin. Oral Implants Res.* 17(5) (2006) 512-520.
- [19] M. Quirynen, R. Vogels, W. Peeters, D. van Steenberghe, I. Naert, A. Haffajee, Dynamics of initial subgingival colonization of ‘pristine’ peri-implant pockets. *Clin. Oral Implants Res.* 17(1) (2006) 25-37.
- [20] W. Heuer, M. Stiesch, W.R. Abraham, Microbial diversity of supra- and subgingival biofilms on freshly

- colonized titanium implant abutments in the human mouth. *Eur. J. Clin. Microbiol. Infect. Dis.* 30(2) (2011) 193-200.
- [21] D. Bosshardt, V. Chappuis, D. Buser, Osseointegration of titanium, titanium alloy and zirconia dental implants: current knowledge and open questions. *Periodontology* 2000. 73 (2017) 22-40.
- [22] B. Zhao, H.C. van der Mei, G. Subbiahdoss, J. de Vries, M. Rustema-Abbing, R. Kuijter, H.J. Busscher, Y. Ren, Soft tissue integration versus early biofilm formation on different dental implant materials. *Dent. Mater.* 30(7) (2014) 716-727.
- [23] L. Rimondini, L. Cerroni, A. Carrassi, P. Torricelli, Bacterial colonization of zirconia ceramic surfaces: an in vitro and in vivo study. *Int. J. Oral Maxillofac. Implants.* 17(6) (2002) 793-798.
- [24] E.D. Avila, M.J. Avila-Campos, C.E. Vergani, D.M. Spolidório, F.A.Jr. Mollo, Structural and quantitative analysis of a mature anaerobic biofilm on different implant abutment surfaces. *J. Prosthet. Dent.* 115(4) (2016) 428-436.
- [25] T. Wassmann, S. Kreis, M. Behr, R. Buegers, The influence of surface texture and wettability on initial bacterial adhesion on titanium and zirconium oxide dental implants. *Int. J. Implant Dent.* 3 (2017) 32. DOI 10.1186/s40729-017-0093-3.
- [26] S. Koka, M.E. Razzoog, T.J. Bloem, S. Syed, Microbial colonization of dental implants in partially edentulous subjects. *J. Prosthet. Dent.* 70(2) (1993) 141-144.
- [27] L.G. Persson, U. Lekholm, A. Leonhardt, G. Dahlén, J. Lindhe, Bacterial colonization on internal surfaces of Branemark system implant components. *Clin. Oral Implants Res.* 7(2) (1996) 90-95.
- [28] A.J. van Winkelhoff, E.G. Winkel, Systemic antibiotic therapy in severe periodontitis. *Curr. Opin. Periodontol.* 4 (1997) 35-40.
- [29] E.G. Salvi, M.M. Furst, N.P. Lang, G.R. Persson One-year bacterial colonization pattern of *Staphylococcus aureus* and other bacteria at implants and adjacent teeth. *Clin. Oral Implants Res.* 19(3) (2008) 242-248.
- [30] M. Quirynen, R. Vogels, M. Pauwels, A.D. Haffajee, S.S. Socransky, N.G. Uzel, D. van Steenberghe, Initial subgingival colonization of 'pristine' pockets. *J. Dent. Res.* 84(4) (2005) 340-344.
- [31] A.L. De Boever, J.A. De Boever, Early colonization of non-submerged dental implants in patients with a history of advanced aggressive periodontitis. *Clin. Oral Implants Res.* 17(1) (2006) 8-17.
- [32] K. Heydenrijk, H.J. Meijer, W.A. van der Reijden, G.M. Raghoobar, A. Vissink, B. Stegenga, Microbiota around root-form endosseous implants: a review of the literature. *Int. J. Oral Maxillofac. Implants.* 17(6) (2002) 829-838.
- [33] A. Leonhardt, C. Bergstrom, U. Lekholm, Microbiologic diagnostics at titanium implants. *Clin. Implant Dent. Relat. Res.* 5(4) (2003) 226-232.
- [34] J. Gerber, D. Wenaweser, L. Heitz-Mayfield, N.P. Lang, G.R. Persson, Comparison of bacterial plaque samples from titanium implant and tooth surfaces by different methods. *Clin. Oral Implants Res.* 17(1) (2006) 1-7.
- [35] W. Keller, U. Bragger, A. Mombelli Peri-implant microflora of implants with cemented and screw retained suprastructures. *Clin. Oral Implants Res.* 9(4) (1998) 209-217.
- [36] M.J. Flynn, J. Slots, Beta-hemolytic streptococci in advanced periodontitis. *Oral Microbiol. Immunol.* 8(5) (1993) 295-297.
- [37] G.E. Romanos, M.T. Bittucci, A. Kokaras, B.J. Paster, Bacterial composition at the implant-abutment connection under loading in vivo. *Clin. Implant Dent. Relat. Res.* 18(1) (2016) 138-145.
- [38] K. Subramani, R.E. Jung, A. Molenberg, C.H. Hammerle, Biofilm on dental implants: a review of the literature. *Int. J. Oral Maxillofac. Implants.* 24(4) (2009) 616-626.
- [39] A. Revathi, A.D. Borrás, A.I. Muñoz, C. Richard, G. Manivasagam, Degradation mechanisms and future challenges of titanium and its alloys for dental implant applications in oral environment. *Mat. Sci. Eng. C. Mater. Biol. Appl.* 76 (2017) 1354-1368.
- [40] L. Baldwin, J.A. Hunt, Host inflammatory response to NiCr, CoCr, and Ti in a soft tissue implantation model. *J. Biomed. Mater. Res. A.* 79(3) (2006) 574-581.
- [41] F. Yu, O. Addison, S.J. Baker, A.J. Davenport, Lipopolysaccharide inhibits or accelerates biomedical titanium corrosion depending on environmental acidity. *Int. J. Oral Sci.* 7(3) (2015) 179-186.
- [42] O. Addison, A.J. Davenport, R.J. Newport, S. Kalra, M. Monir, J.F. Mosselmans, D. Proops, R.A. Martin, Do 'passive' medical titanium surfaces deteriorate in service in the absence of wear? *J. R. Soc. Interface.* 9(76) (2012) 3161-3164.
- [43] P. Hallam, F. Haddad, J. Cobb, Pain in the well-fixed, aseptic titanium hip replacement. The role of corrosion. *J. Bone Joint Surg. Br.* 86(1) (2004) 27-30.
- [44] S. Virtanen, I. Milosev, E. Gomez-Barrena, R. Trebse, J. Salo, Y.T. Kontinen, Special modes of corrosion under physiological and simulated physiological conditions. *Acta Biomater.* 4(3) (2008) 468-476.
- [45] J.C. Souza, P. Ponthiaux, M. Henriques, R. Oliveira, W. Teughels, J.P. Celis, L.A. Rocha, Corrosion behaviour of titanium in the presence of *Streptococcus mutans*. *J. Dent.* 41(6) (2013) 528-534.
- [46] M.N. Oliveira, W.V.H. Schunemann, M.T. Mathew, B. Henriques, R.S. Magini, W. Teughels, J.C.M. Souza, Can degradation products released from dental implants affect peri-implant tissues? *J. Periodont. Res.* 53 (2018) 1-11.
- [47] M.G. Newman, T.F. Flemming, Periodontal considerations of implants and implant associated microbiota. *J. Dent. Educ.* 52(12) (1988) 737-744.
- [48] C. Nascimento, N. Monesi, I.Y. Ito, J.P. Issa, R.F. de Albuquerque Junior, Bacterial diversity of periodontal and implant-related sites detected by the DNA Checkerboard method. *Eur. J. Clin. Microbiol. Infect. Dis.* 30(12) (2011) 1607-1613.
- [49] S. Renvert, G.R. Persson, Periodontitis as a potential risk factor for peri-implantitis. *J. Clin. Periodontol.* 36(Suppl 10) (2009) 9-14.
- [50] J. Neilands, C. Wickstrom, B. Kinnby, J.R. Davies, J. Hall, B. Friberg, G. Svensater, Bacterial profiles and proteolytic activity I peri-implantitis versus healthy sites. *Anaerobe.* 35(Pt A) (2015) 28-34.
- [51] T. Koyanagi, M. Sakamoto, Y. Takeuchi, N. Maruyama, M. Ohkuma, Y. Izumi, Comprehensive microbiological

- findings in peri-implantitis and periodontitis. *J. Clin. Periodontol.* 40(3) (2013) 218-226.
- [52] J.A. Shibli, M.C. Martins, R.F. Lotufo, E.Jr. Marcantonio Microbiologic and radiographic analysis of ligature-induced peri-implantitis with different dental implant surfaces. *Int. J. Oral Maxillofac. Implants.* 18(3) (2003) 383-390.
- [53] K. Heydenrijk, H.J. Meijer, W.A. van der Reijden, G.M. Raghoobar, A. Vissink, B. Stegenga, Microbiota around root-form endosseous implants: a review of the literature. *Int. J. Oral Maxillofac. Implants.* 17(6) (2002) 829-838.
- [54] T. van de Velde, E. Thevissen, G.R. Persson, C. Johansson, H. De Bruyn, Two-year outcome with Nobel Direct implants: a retrospective radiographic and microbiologic study in 10 patients. *Clin. Implant Den. Rel. Res.* 11(3) (2009) 183-193.
- [55] D. Kohavi, R. Greenberg, E. Raviv, M.N. Sela, Subgingival and supragingival microbial flora around healthy osseointegrated implants in partially edentulous patients. *Int. J. Oral Maxillofac. Implants.* 9(6) (1994) 673-678.
- [56] M. Quirynen, N. van Assche, Microbial changes after full-mouth tooth extraction, followed by 2-stage implant placement. *J. Clin. Periodontol.* 38(6) (2011) 581-589.
- [57] S.L. Devides, A.T. Franco, Evaluation of peri-implant microbiota using the polymerase chain reaction in completely edentulous patients before and after placement of implant-supported prostheses submitted to immediate load. *Int. J. Oral Maxillofac. Implants.* 21(2) (2006) 262-269.
- [58] A. Sachdeo, A.D. Haffajee, S.S. Socransky, Biofilms in the edentulous oral cavity. *J. Prosthodont.* 17(5) (2008) 348-356.
- [59] J.A. Shibli, L. Melo, D.S. Ferrari, L.C. Figueiredo, M. Faveri, M. Feres, Composition of supra- and subgingival biofilm of subjects with healthy and diseased implants. *Clin. Oral Implants Res.* 19(10) (2008) 975-982.
- [60] A.D. Haffajee, S.S. Socransky, Microbial etiological agents of destructive periodontal diseases. *Periodontol.* 2000. 5 (1994) 78-111.
- [61] S.S. Socransky, A.D. Haffajee, M.A. Cugini, C. Smith, R.L.Jr. Kent, Microbial complexes in subgingival plaque. *J. Clin. Periodontol.* 25(2) (1998) 134-144.
- [62] J. Slots, T.E. Rams, D. Feik, H.D. Taveras, G.M. Gillespie, Subgingival microflora of advanced periodontitis in the Dominican Republic. *J. Periodontol.* 62(9) (1991) 543-547.
- [63] L.F. Zhuang, R.M. Watt, N. Matheos, M.S. Si, H.C. Lai, N.P. Lang, Periodontal and peri-implant microbiota in patients with healthy and inflamed periodontal and peri-implant tissues. *Clin. Oral Implants Res.* 27(1) (2016) 13-21.
- [64] M. Morra, C. Cassinelli, D. Bollati, G. Cascardo, M. Bellanda, Adherent endotoxin on dental implant surfaces: A reappraisal. *J. Oral Implantol.* 41(1) (2015) 10-16.
- [65] E.S. Rosenberg, J.P. Torosian, J. Slots, Microbial differences in 2 clinically distinct types of failures of osseointegrated implants. *Clin. Oral Implants Res.* 2(3) (1991) 135-144.
- [66] A.D. Pye, D.E. Lockhart, M.P. Dawson, C.A. Murray, A.J. Smith, A review of dental implants and infection. *J. Hosp. Infect.* 72(2) (2009) 104-110.
- [67] D. Apatzidou, D.F. Lappin, G. Hamilton, C.A. Papadopoulos, A. Konstantinidis, M.P. Riggio, Microbiome of peri-implantitis affected and healthy dental sites in patients with a history of chronic periodontitis. *Arch. Oral Biol.* 83 (2017) 145-152.
- [68] M. Quirynen, M.A. Listgarten, The distribution of bacterial morphotypes around natural teeth and titanium implants ad modum Branemark. *Clin. Oral Implants Res.* 1(1) (1990) 8-12.
- [69] M.M. Furst, G.E. Salvi, N.P. Lang, G.R. Persson, Bacterial colonization immediately after installation on oral titanium implants. *Clin. Oral Impl. Res.* 18(4) (2007) 501-508.
- [70] M. Esposito, M.G. Grusovin, H.V. Worthington, Interventions for replacing missing teeth: antibiotics at dental implant placement to prevent complications. *Cochrane Database Syst. Rev.* 7 (2013) CD004152.
- [71] E. Romeo, M. Ghisolfi, D. Carmagnola, Peri-implant diseases. A systematic review of the literature. *Minerva Stomatol.* 53(5) (2004) 215-230.
- [72] R.P. Teles, M. Patel, S.S. Socransky, A.D. Haffajee, Disease progression in periodontally healthy and maintenance subjects. *J. Periodontol.* 79(5) (2008) 784-794.
- [73] L. Meijndert, W.A. van der Reijden, G.M. Raghoobar, H.J. Meijer, A. Vissink, Microbiota around teeth and dental implants in periodontally healthy, partially edentulous patients: is pre-implant microbiological testing relevant? *Eur. J. Oral Sci.* 118(4) (2010) 357-363.
- [74] A. Mombelli, Microbial colonization of the periodontal pocket and its significance for periodontal therapy. *Periodontology 2000.* 76 (2018) 85-96.
- [75] M. Gross, I. Abramovich, E.I. Weiss, Microleakage at the abutment-implant interface of osseointegrated implants: a comparative study. *Int. J. Oral Maxillofac. Implants.* 14(1) (1999) 94-100.
- [76] T. Handal, I. Olsen, C.B. Walker, D.A. Caugant, Beta-lactamase production and antimicrobial susceptibility of subgingival bacteria from refractory periodontitis. *Oral Microbiol. Immunol.* 19(5) (2004) 303-308.
- [77] V.K. Jansen, G. Conrads, E.J. Richter, Microbial leakage and marginal fit of the implant-abutment interface. *Int. J. Oral Maxillofac. Implants.* 12(4) (1997) 527-540.
- [78] M. Tallarico, L. Canullo, M. Caneva, M. Ozcan, Microbial colonization at the implant-abutment interface and its possible influence on periimplantitis: A systematic review and meta-analysis. *J. Prosthodont. Res.* 61 (2017) 233-241.
- [79] J.S. Guindy, C.E. Besimo, R. Besimo, H. Schiel, J. Meyer, Bacterial leakage into and from prefabricated screw-retained implant-borne crowns in vitro. *J. Oral Rehabil.* 25(6) (1998) 403-408.
- [80] L. Steinebrunner, S. Wolfart, K. Bossmann, M. Kern, In vitro evaluation of bacterial leakage along the implant-abutment interface of different implant designs. *Int. J. Oral Maxillofac. Implants.* 20(6) (2005) 875-881.
- [81] M. Quirynen, D. van Steenberghe, Bacterial colonization of the internal part of two-stage implants.

- An in vivo study. Clin. Oral Implants Res. 4(3) (1993) 158-161.
- [82] C.E. Besimo, J.S. Guindy, D. Lewetag, J. Meyer, Prevention of bacterial leakage into and from prefabricated screw-retained crowns on implants in vitro. Int. J. Oral Maxillofac. Implants. 14(5) (1999) 654-660.
- [83] G. Orsini, B. Assenza, A. Scarano, M. Piattelli, A. Piattelli, Surface analysis of machined versus sandblasted and acid-etched titanium implants. Int. J. Oral Maxillofac. Implants. 15(6) (2000) 779-784.
- [84] A. Piattelli, A. Scarano, M. Paolantonio, B. Assenza, G.C. Leghissa, G. Di Bonaventura, G. Catamo, R. Piccolomini, Fluids and microbial penetration in the internal part of cement-retained versus screw-retained implant-abutment connections. J. Periodontol. 72(9) (2001) 1146-1150.
- [85] L. Rimondini, C. Marin, F. Brunella, M. Fini, Internal contamination of a 2-component implant system after occlusal loading and provisionally luted reconstruction with or without a washer device. J. Periodontol. 72(12) (2001) 1652-1657.
- [86] T. Koutouzis, S. Wallet, N. Calderon, T. Lundgren, Bacterial colonization of the implant-abutment interface using an in vitro dynamic loading model. J. Periodontol. 82(4) (2011) 613-618.
- [87] M. Tesmer, S. Wallet, T. Koutouzis, T. Lundgren, Bacterial colonization of the dental implant fixture-abutment interface: an in vitro study. J. Periodontol. 80(12) (2009) 1991-1997.
- [88] F. Grecchi, M. di Girolamo, F. Cura, V. Candotto, F. Carinci, A new system of implant abutment connection: how to improve a two piece implant system sealing. Oral & Implantology. 10(3) (2017) 234-240.